

REGEN-COV (Casirivimab with Imdevimab) Infusion Order Form

Patient Name: _____ DOB: _____ Phone: _____
Address: _____ City: _____ State: _____ Zip: _____
Patient SS #: _____ Allergies: _____ Pt. Weight: _____ lbs/kg
Physician: _____ NPI: _____
Insurance Name: _____ Patient ID: _____
Please Circle: Date of First Symptom or Exposure Onset: _____ COVID Positive Date: _____

Please send face sheet or copy of insurance cards. If Medicare patient, please include SSN

Patient Eligibility

Exclusion Criteria: Patients meeting any of the following criteria are NOT ELIGIBLE for Casirivimab with Imdevimab therapy

- a. who are hospitalized due to COVID-19
- b. who require oxygen therapy due to COVID-19
- c. who require an increase in baseline oxygen flow rate due to COVID-19 in those on chronic oxygen therapy due to underlying non-COVID-19 related comorbidity

Inclusion Criteria: The following criteria are required to qualify a patient for Casirivimab with Imdevimab therapy

Check all that apply:

- Patient is 12 years of age or older weighing at least 40 kg who are at high risk (see criteria below) for progression to severe COVID-19, including hospitalization or death, and are not fully vaccinated **or** who are not expected to mount an adequate immune response to complete SARS-CoV-2 vaccination (for example, individuals with immunocompromising conditions including those taking immunosuppressive medications) **and**

Prophylaxis Patients only:

- have been exposed to an individual infected with SARS-CoV-2 consistent with close contact criteria per Centers for Disease Control and Prevention (CDC) **or**
- who are at high risk of exposure to an individual infected with SARS-CoV-2 because of occurrence of SARS-CoV-2 infection in other individuals in the same institutional setting (for example, nursing homes, prisons)

COVID Positive Patients: Therapy must begin within 10 days of Symptom onset regardless of COVID positive test date

High Risk Patients must have at least one of the following (select all that apply):

- Older age (for example, age ≥ 65 years of age)
- Obesity or being overweight (for example, BMI > 25 kg/m², or if age 12-17, have BMI ≥ 85 th percentile for their age and gender based on CDC growth charts, https://www.cdc.gov/growthcharts/clinical_charts.htm)
- Pregnancy
- Chronic kidney disease
- Diabetes
- Immunosuppressive disease or immunosuppressive treatment
- Cardiovascular disease (including congenital heart disease) or hypertension
- Chronic lung diseases (for example, chronic obstructive pulmonary disease, asthma [moderate-to-severe], interstitial lung disease, cystic fibrosis and pulmonary hypertension)
- Sickle cell disease
- Neurodevelopmental disorders (for example, cerebral palsy) or other conditions that confer medical complexity (for example, genetic or metabolic syndromes and severe congenital anomalies)
- Having a medical-related technological dependence (for example, tracheostomy, gastrostomy, or positive pressure ventilation (not related to COVID-19))
- Other medical conditions or factors that may place individual patients at high risk for severe COVID-19:

Physician Signature: _____ Printed: _____ Date: _____

IV Infusion Orders

- Casirivimab 600mg with Imdevimab 600mg in 100 mL 0.9% Sodium Chloride** to be infused IV via gravity or infusion pump over 30 minutes x 1 dose
(Must use a 0.2 or 0.22 micron filter for administration)
- Subsequent Repeat Dosing:** For patients in whom repeat dosing is determined to be appropriate for ongoing exposure to SARS-CoV-2 for longer than 4 weeks and who are not expected to mount an adequate immune response to complete SARS-CoV-2 vaccination
- The initial intravenous infusion may be followed by subsequent repeat dosing of 300 mg of Casirivimab and 300 mg of Imdevimab by IV infusion once every 4 weeks for the duration of ongoing exposure.
- 50mL 0.9% Sodium Chloride.** Once infusion is complete, flush the infusion line with 50mL 0.9% Sodium Chloride to ensure delivery of required dose.
- Clinically monitor patients during infusion and observe patients for at least 1 hour after infusion is complete.

Emergency medications for Potential Acute Infusion Reactions

- Anaphylaxis Kit per **Amber Specialty Pharmacy Home infusion anaphylaxis treatment protocol**
- Albuterol Inhaler to be used as needed for severe respiratory reactions
- Solu-Medrol 125mg/2mL IV be used as needed for severe respiratory reactions and/or anaphylactic reactions (e.g. Angioedema) as instructed by Physician.

Anaphylaxis Kit Contents	
Epinephrine 1mg Vial (1:1000 USP) Diphenhydramine HCL (50 mg/1mL vial) 0.9% Sodium Chloride (500 mL) 2 x 1mL syringe w/ 25G x 1" needle	2 x 3mL syringe w/ 25G x 1.5" needle, Non-vented IV Set Alcohol wipes

Vascular Access Device (VAD) Orders:

Peripheral Vascular Access Device: Skilled nursing to assess and insert peripheral access device for administration of Casirivimab with Imdevimab.

Other: _____

Other: _____

Clinical Services:

Pharmacy Services:

Assessment of patient eligibility, administration method, education on medication side effects, interactions, adverse reactions, and infusion-related reactions.

Nursing Services:

Skilled nursing to administer Casirivimab with Imdevimab, patient assessment, and monitoring.

Physician Signature: _____	Printed: _____	Date: _____
----------------------------	----------------	-------------

Monitoring

- Document Vital Signs: Temperature, HR, RR, Pulse Ox taken before medication initiation; immediately after medication administration; and 1 hour post medication administration
- Medical professional to monitor patient 1-hour post medication administration
- Document time of medication administration
- Note any adverse reactions

Vital Sign	Prior to Med Administration	Immediately after Med Administration	1 Hour Post Medication Administration
Temp			
HR			
RR			